Ce501-503

# T.J. KOBAYASHI LAB.

# [Mathematics & Informatics for Life Science]

#### Laboratory for Quantitative Biology

Systems biology, Quantitative biology, Theoretical Biology

Department of Mathematical informatics

Department of Electrical Engineering and Information Systems (EEIS)

**Frontier of Biology as an Interdisciplinary Science** 

Cellular phenomena such as development, immunity, and







cancer are intrinsically dynamics. Recent development of bio-imaging technology enables us to directly and quantitatively observe these dynamics. This innovation in turn drives the frontier of biology to become an interdisciplinary field by incorporating the knowledge and 1.6 technologies of other fields such as engineering, informatics,<sup>1.25</sup> and mathematics where quantitative data and analysis are 0.75 commonly used.

### **Information Technology for Biology**

Information technology such as image and data analysis is indispensable to exploit the potential information that bio-imaging data have. We have developed methods for 4D tracking of mouse embryogenesis (Fig.1-2) from 4D bioimaging data, and prediction of peptide folding and its determinants (Fig. 3). By using these methods, we have revealed that desynchronization of circadian cells underlies singularity phenomena in which stop of circadian oscillation is induced by unusual mid-night light.



Ukai, Kobayashi, et al., (2007), Nat. Cell Biol.



Fig. 3 Potential energy PCA analysis of peptide folding



## Theory for robust operation of a cell with noisy components

Mathematical modeling is also crucial to design principles of dynamic (A) understand biological systems. For example, bio-imaging revealed that cellular systems can operate robustly even with substantial noise in their components. However, its mechanism still (C) remains unknown. By combining statistical theory with that for stochastic chemical reaction, we have theoretically clarified that a very simple intracellular reaction can conduct surprising <sup>(E)</sup> information-processing to extract relevant information from very noisy signal. This theory may also be employed to design noise-immune





